## MATH 223: Axioms.

## Field Axioms

A field is a set of elements  $\mathbf{F}$  which we call *scalars* when used in a vector space. A quick check verifies that the real numbers  $\mathbf{R}$ , the complex numbers  $\mathbf{C}$  and the rational numbers  $\mathbf{Q}$  all are examples of fields. The integers  $\mathbf{Z}$  are not (because no multiplicative inverses). There are other useful examples of fields which we do not use in this course.

In what follows  $\alpha, \beta, \gamma$  are arbitrary elements of **F** 

$$\forall \alpha, \beta \in \mathbf{F} \ \alpha + \beta = \beta + \alpha \text{ (commutativity)}$$

$$\forall \alpha, \beta, \gamma \in \mathbf{F} \ \alpha + (\beta + \gamma) = (\alpha + \beta) + \gamma \text{ (associativity)}$$

$$\exists 0 \text{ with } \forall \alpha \in \mathbf{F} \ \alpha + 0 = \alpha$$

$$\forall \alpha \in F \ \exists - \alpha \in F \text{ with } \alpha + (-\alpha) = 0$$

$$\forall \alpha, \beta \in \mathbf{F} \ \alpha \beta = \beta \alpha \text{ (commutativity)}$$

$$\forall \alpha, \beta, \gamma \in \mathbf{F} \ \alpha (\beta \gamma) = (\alpha \beta) \gamma \text{ (associativity)}$$

$$\exists 1 \text{ with } \forall \alpha \in \mathbf{F} \ \alpha 1 = \alpha$$

$$\forall \alpha \in \mathbf{F} \text{ with } \alpha \neq 0, \exists \alpha^{-1} \text{ with } \alpha \alpha^{-1} = 1$$

$$\forall \alpha, \beta, \gamma \in \mathbf{F} \ \alpha (\beta + \gamma) = \alpha \beta + \alpha \gamma$$

## Vector Spaces Axioms

We have as set V of *vectors* and a field  $\mathbf{F}$  (typically  $\mathbf{R}$  in this course) of scalars.

In what follows  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  are arbitrary elements of V and  $k, \ell$  are arbitrary elements of  $\mathbf{F}$ .

$$\forall \mathbf{u}, \mathbf{v} \in V, \mathbf{u} + \mathbf{v} \in V \text{ (closure)}$$

$$\forall \mathbf{u} \in V \ \forall k \in \mathbf{F}, k\mathbf{u} \in V \text{ (closure)}$$

$$\forall \mathbf{u}, \mathbf{v} \in V, \mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u} \text{ (commutativity)}$$

$$\forall \mathbf{u}, \mathbf{v}, \mathbf{w} \in V, (\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w}) \text{ (associativity)}$$

$$\exists \mathbf{0} \text{ (that is unique) with } \forall \mathbf{u} \in V, \mathbf{0} + \mathbf{v} = \mathbf{v}$$

$$\forall k \in \mathbf{F} \ \forall \mathbf{u}, \mathbf{v} \in V \ k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v} \text{ (distributivity)}$$

$$\forall k, \ell \in \mathbf{F} \ \forall \mathbf{u} \in V \ (k + \ell)\mathbf{u} = k\mathbf{u} + \ell\mathbf{u} \text{ (distributivity)}$$

$$\forall k, \ell \in \mathbf{F} \ \forall \mathbf{u} \in V \ k\ell\mathbf{u} = k(\ell\mathbf{u})$$

$$\forall \mathbf{u} \in V \ 1 \cdot \mathbf{u} = \mathbf{u}$$