

- Let $A \in \mathbf{R}^{m \times n}$ have rank 1. Show that there exist non-zero vectors $x \in \mathbf{R}^m$ and $y \in \mathbf{R}^n$ so that $A = xy^T$. (Hint: Try a simple case and also compute xy^T for some simple choices for x and y .) (Comment: You could explore how to generalize such a result to higher rank.)
- Determine bases for the following subspaces of \mathbf{R}^3 .
 - the line $x = 5t, y = -2t, z = t$.
 - all vectors of the form $(a, b, c)^T$ such that $a - 3b = 2c$.
- Let

$$A = \begin{bmatrix} 0 & 1 & 1 & 2 & -3 & 1 \\ 0 & 2 & 0 & 6 & -6 & 0 \\ 0 & 3 & 7 & 2 & -9 & 7 \\ 0 & 2 & 2 & 4 & -4 & 3 \end{bmatrix}$$

Determine a basis for the column space of A (chosen from columns of A) and determine a basis for the row space of A . Also give a basis for the *nullspace* of A , namely $\{\mathbf{x} \in \mathbf{R}^6 : A\mathbf{x} = \mathbf{0}\}$.

- Show that the set of all vectors $(b_1, b_2, b_3, b_4)^T$ such that the system below is consistent (i.e. can be solved)

$$\begin{bmatrix} 2 & 3 & 1 \\ 4 & 3 & 3 \\ 1 & 3 & 0 \\ 2 & 0 & 2 \end{bmatrix} \mathbf{x} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}$$

is a subspace of \mathbf{R}^4 . Then find a basis of the subspace.

- Let A be an $n \times n$ matrix with various eigenvalues including λ and μ with $\lambda \neq \mu$. Let L, M be the eigenspaces associated with eigenvalues λ, μ respectively. (That is, L is the set of all eigenvectors with eigenvalue λ ; M is the set of all eigenvectors with eigenvalue μ .) Let $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_p\}$ be a basis for L and let $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_q\}$ be a basis for M . Show that $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_p, \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_q\}$ is a linearly independent set of $p + q$ vectors. (Hint: try $p = 1$ and $q = 1$ to start). (Comment: You could explore the case if there were three different eigenvalues and three bases for the eigenspaces).
- Let $\mathbf{R}^{n \times n}$ denote the vector space of all $n \times n$ matrices (over \mathbf{R}). Consider following transformation $f : \mathbf{R}^{n \times n} \rightarrow \mathbf{R}^{n \times n}$

$$f(A) = A^T.$$

Show that this is a linear transformation.

We say that a matrix A is *symmetric* if $A^T = A$ and we say that a matrix A is *skew-symmetric* if $A^T = -A$.

- Warmup question: Give a basis for $\mathbf{R}^{n \times n}$. How many elements are in your basis?
- What is the dimension of the eigenspace of eigenvalue 1 for f ? Explain.
- What is the dimension of the eigenspace of eigenvalue -1 for f ? Explain.
- Now use the previous question (and other facts) to show that any $A \in \mathbf{R}^{n \times n}$ is a linear combination of a symmetric matrix and a skew-symmetric matrix (you could show this directly of course but I'm asking you to use linear independence/dimension arguments).