

1. Solve the differential equation given the initial conditions  $x_1(0) = -3$ ,  $x_2(0) = 4$ .

$$\begin{aligned}\frac{d}{dt}x_1(t) &= \quad \quad + x_2(t) \\ \frac{d}{dt}x_2(t) &= -2x_1(t) - 2x_2(t)\end{aligned}$$

2. You are given a 3 dimensional vector space  $V \subseteq \mathbf{R}^5$ . Could there be a  $3 \times 6$  matrix  $A$  with nullspace of  $A$  being  $V$ ? Explain. Could there be  $6 \times 5$  matrix  $B$  with nullspace of  $B$  being  $V$ ? Explain. In either case, if you were given a basis for the three dimensional space  $V$ , how would you find the desired matrix assuming it exists.

3. Consider the two planes  $\pi_1: x - y + 2z = 3$  and  $\pi_2: x + 2y + 3z = 6$ .

a) Find the intersection of  $\pi_1$  and  $\pi_2$  in vector parametric form.

b) What is the angle (or just the cosine of the angle) formed by the two planes? (This is defined as the angle between their normal vectors. A normal vector is a vector orthogonal  $u - v$  for every  $u, v$  in the plane.)

c) Find the distance of the point  $(-1, 2, 2)$  to the plane  $\pi_1$ . (That is, find the distance between  $(-1, 2, 2)$  and the closest point in  $\pi_1$ .)

d) Find the equation of the plane parallel to  $\pi_1$  through the point  $(3, 2, 0)$ .

e) Imagine the direction  $(0, 0, 1)^T$  as pointing straight up from your current position  $(0, 0, 0)^T$  in 3-space and the plane  $\pi_2$  as a physical plane. If a marble is placed on  $\pi_2$  at the point  $(6, 0, 0)^T$ , what direction will the marble roll under the influence of gravity?

4. Given a matrix  $A \in \mathbf{R}^{n \times n}$ , we define the trace

$$\text{tr}(A) = \sum_{i=1}^n A_{i,i},$$

i.e., the sum of the diagonal. This is an important quantity.

a) Let  $A, B \in \mathbf{R}^{n \times n}$ . Show that  $\text{tr}(AB) = \text{tr}(BA)$ . *Hint: You may wish to express  $AB$  using the dot products between rows of  $A$  and columns of  $B$ . To be precise, let  $u_1, u_2, \dots, u_n$  be the columns of  $A^T$  (rows of  $A$ ) and  $v_1, v_2, \dots, v_n$  be the columns of  $B$ . Then  $(AB)_{i,j} = u_i \cdot v_j$ . You can then show that  $\text{tr}(AB)$  is the dot product between  $A^T$  and  $B$  (it's up to you to define this dot product between matrices).*

b) Suppose that  $A$  can be diagonalized as  $A = MDM^{-1}$  where  $D$  is a diagonal matrix of eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ . Show that

$$\text{tr}(A) = \sum_{i=1}^n \lambda_i.$$

**Important note:** *The above equality is true even if  $A$  cannot be diagonalized. In other words, let  $\lambda_1, \lambda_2, \dots, \lambda_n$  be the  $n$  solutions to the characteristic equation  $\det(A - \lambda I) = 0$ . By the Fundamental Theorem of Algebra, there are always  $n$  solutions when counted with multiplicity. These are the eigenvalues of  $A$ . Then*

$$\text{tr}(A) = \sum_{i=1}^n \lambda_i.$$

*You may use this fact without proof.*

5. Let  $A$  be a  $n \times n$  matrix of real entries satisfying  $A^2 = -I$ . Show that
- $A$  is invertible (or *nonsingular*)
  - $A$  has no real eigenvalues
  - $n$  is even
  - (harder question)  $\det(A) = 1$ . (Hint: Try using the previous question.)
6. Consider two vector spaces  $U, V$ , subspaces of  $\mathbb{R}^m$ . Define  $U + V = \{\mathbf{u} + \mathbf{v} : \mathbf{u} \in U, \mathbf{v} \in V\}$ . (This is called the Minkowski sum.) Show that  $U + V$  is a vector space. Now show that  $\dim(U) + \dim(V) = \dim(U \cap V) + \dim(U + V)$ .
- (Hint: if we have an  $m \times n$  matrix  $A$  then  $n = \dim(\text{nullspace}(A)) + \text{rank}(A)$ . How should we form  $A$ ? )